

Ascorbic Acid (AsA) and Total Ascorbic Acid (T-AsA) Content Assay Kit

Note: Take two or three different samples for prediction before test.

Detection equipment: Spectrophotometer

Cat No: NA0252

Size: 50T/24S

Components:

Extract Solution: 40 mL×1, storage at 4°C.

Reagent I: Powder×1, store at -20°C. Add 3.3 mL of distilled water when the solution will be used. The unused reagents can be stored at -20°C and avoid light.

Reagent II: Liquid 20 mL×1, store at 4°C.

Reagent III: Liquid 3 mL×1, store at 4°C.

Reagent IV: Liquid 30 mL×1, store at 4°C.

Reagent V: Liquid 20 mL×1, store at 4°C.

Reagent VI: Powder×1, store at 4°C. Add 12 mL of 70% ethanol(v/v) before use, mix thoroughly.

Reagent VII: Liquid 12 mL×1, store at 4°C.

Standard: Powder×1, store at 4°C. Add 1.136 mL of extract solution before use, mix thoroughly. Add 0.99 mL distilled water to 0.01 mL of above standard solution, mix thoroughly and to be prepared as 500 nmol/mL AsA standard solution for use.

Description:

AsA is also called Vitamin C. AsA is the substrate of coenzyme, free radical scavenger, electron copolymer/receptor, biosynthesis of oxalate and tartrate. As the most important antioxidant in plant cells, AsA has important function in protecting chloroplast from oxidizing. It is also one of the important indexes to measure the quality of crop products. DHA is a reversible oxidized form of AsA. It forms a redox system with ascorbic acid in the living body and has the function of an electron acceptor.

ASA has reductive ability, which can reduce Fe^{3+} to Fe^{2+} . Fe^{2+} and 2,2'-bipyridine form a pink complex, with a characteristic absorption peak at 525 nm. DTT can reduce DHA to generate ASA, for this can be used to detect the total ascorbic acid(ASA + DHA) content of samples.

Required but not provided

Mortar, ice, low temperature centrifuge, spectrophotometer, 1 mL glass cuvette, transferpettor, ethanol and distilled water.

Protocol:

I. Sample extraction:

1. Tissue:

Accordance the ratio of tissue(g) : extract solution volume (mL)=1: 5~10, (add 1 mL of extract solution to 0.1 g of tissue), homogenate on ice. Centrifuge at 13000 ×g for 10 minutes at 4°C, take the supernatant and place it on ice for testing.

2. Bacteria or cells:

Accordance the ratio of cells amount(10^4) : extract solution volume (mL)=500~1000: 1, (add 1 mL of extract solution to 5 million cells). Ultrasonic on ice bath to smash cells, (powder 300w, ultrasonic 3s, interval 7s for 3 minutes). Centrifuge at 13000 ×g for 10 minutes at 4°C, take the supernatant and place it on ice for testing.

3. Serum:

Add 500 μL of extract solution to 500 μL of sample, vortex mixing. Centrifuge at 13000 ×g for 10 minutes at 4°C, take the supernatant and place it on ice for testing.

II. Determination procedure

1. Preheat ultraviolet spectrophotometer for 30 minutes, adjust wavelength to 525 nm, set zero with distilled water.

2. Determination of ASA content:

Add reagents with the following list:

Reagent (μL)	Test tube (T)	Contrast tube (C)	Blank tube 1 (B1)	Blank tube 2 (B2)	Standard tube (S)
Sample	50	50	-		-
Extract solution	-	-	50	50	-
Standard solution	-	-			50
Reagent II	200	200	200	200	200
Reagent IV	250	250	250	250	250
Reagent V	200	200	200	200	200
Reagent VI	200	-	200	-	200
70% ethanol	-	200	-	200	-
Reagent VII	100	100	100	100	100

Mix well, react in water bath at 42°C for 40 minutes, cooling with cold water. Determine the absorption value at 525 nm, record as A_T , A_C , A_{B1} , A_{B2} and A_S . Calculate $\Delta A_{T1}=(A_T - A_C)-(A_{B1}-A_{B2})$, $\Delta A_{S1}=A_S-A_{B1}$.

Note: When adding Reagent VII, put the tips of transferpette under the liquid level, do not drop it in the air, otherwise the liquid will be turbid. The Blank tube 1, Blank tube 2 and Standard tube only needs to be measured one or twice.

3. Determination of ASA content:

Add reagents with the following list:

Reagent (μL)	Test tube (T)	Contrast tube (C)	Blank tube 1 (B1)	Blank tube 2 (B2)	Standard tube (S)
Sample	50	50	-	-	-
Extract solution	-	-	50	50	-

Standard solution	-	-	-	-	50
Reagent I	50	50	50	50	50
Reagent II	100	100	100	100	100
Mix well, react in water bath at 42°C for 15 minutes.					
Reagent III	50	50	50	50	50
Mix well, place at room temperature for 1 minute.					
Reagent IV	250	250	250	250	250
Reagent V	200	200	200	200	200
Reagent VI	200	-	200	-	200
70% ethanol	-	200	-	200	-
Reagent VII	100	100	100	100	100
Mix well, react in water bath at 42°C for 40 minutes, cooling with cold water. Determine the absorption value at 525 nm, record as A _T , A _C , A _{B1} , A _{B2} and A _S . Calculate $\Delta A_{T2}=(A_T - A_C)-(A_{B1}-A_{B2})$, $\Delta A_{S2}=A_S-A_{B1}$.					

Note: When adding Reagent VII, put the tips of transferpette under the liquid level, do not drop it in the air, otherwise the liquid will be turbid. The Blank tube 1, Blank tube 2 and Standard tube only needs to be measured one or twice.

III. Calculation of ASA/T-ASA Content:

A. Calculation of ASA content:

1. Sample weight:

$$AsA(nmol/g) = [C_S \times \Delta A_{T1} \div \Delta A_{S1} \times V_{SV}] \div (W \times V_{SV} \div V_{STV}) = 500 \times \Delta A_{T1} \div \Delta A_{S1} \div W$$

2. Cells:

$$AsA(nmol/10^4 \text{ cell}) = [C_S \times \Delta A_{T1} \div \Delta A_{S1} \times V_{SV}] \div (N \times V_{SV} \div V_{STV}) = 500 \times \Delta A_{T1} \div \Delta A_{S1} \div N$$

3. Liquids:

$$AsA (nmol/mL) = [C_S \times \Delta A_{T1} \div \Delta A_{S1} \times V_{SV}] \times 2 = 1000 \times \Delta A_{T1} \div \Delta A_{S1}$$

C_S: 500 nmol/mL;

V_{STV}: The volume of supernatant after centrifugation, 1 mL;

V_{SV}: The volume of supernatant added into the reaction system, 0.05 mL;

W: Sample weight, g;

V_{Liq}: The volume of sample added during extraction, 0.5 mL;

V_E: The volume of extract solution added during extraction, 0.5 mL;

2: The ratio of dilution, $(V_{Liq} + V_E) / V_{Liq} = (500 \mu L + 500 \mu L) / 500 \mu L = 2$.

N: The number of cells.

B. Calculation of T-ASA content:

$$T-AsA(nmol/g) = [C_S \times \Delta A_{T2} \div \Delta A_{S2} \times V_{SV}] \div (W \times V_{SV} \div V_{STV}) = 500 \times \Delta A_{T2} \div \Delta A_{S2} \div W$$

2. Cells:

$$T-AsA(nmol/10^4 \text{ cell}) = [C_S \times \Delta A_{T2} \div \Delta A_{S2} \times V_{SV}] \div (N \times V_{SV} \div V_{STV}) = 500 \times \Delta A_{T2} \div \Delta A_{S2} \div N$$

3. Liquids

$$T-AsA (nmol/mL) = [C_S \times \Delta A_{T2} \div \Delta A_{S2} \times V_{SV}] \times 2 = 1000 \times \Delta A_{T2} \div \Delta A_{S2}$$

C_S : 500 nmol/mL;

V_{STV} : The volume of supernatant after centrifugation, 1 mL;

V_{SV} : The volume of supernatant added into the reaction system, 0.05 mL;

W : Sample weight, g;

V_{Liq} : The volume of sample added during extraction, 0.5 mL;

V_E : The volume of extract solution added during extraction, 0.5 mL;

2: The ratio of dilution, $(V_{Liq} + V_E)/V_{Liq} = (500 \mu\text{L} + 500 \mu\text{L})/500 \mu\text{L} = 2$.

N : The number of cells.

C. Calculation of DHA content:

1. Sample weight:

$$\text{DHA (nmol/g)} = 500 \times (\Delta A_{T2} \div \Delta A_{S2} - \Delta A_{T1} \div \Delta A_{S1}) \div W$$

2. Cells:

$$\text{DHA (nmol/10}^4 \text{ cell)} = 500 \times (\Delta A_{T2} \div \Delta A_{S2} - \Delta A_{T1} \div \Delta A_{S1}) \div N$$

3. Liquids:

$$\text{DHA (nmol/mL)} = 1000 \times (\Delta A_{T2} \div \Delta A_{S2} - \Delta A_{T1} \div \Delta A_{S1})$$

W : Sample weight, g;

N : The number of cells.

Note:

1. When adding Reagent VII, put the tips of transferpette under the liquid level, do not drop it in the air, otherwise the liquid will be turbid.
2. The Standard tube only needs to be measured one or twice.
3. When the measured absorption value is greater than 1, it is recommended to measure after dilution with extract solution, multiply the corresponding dilution ratio in calculation.
4. This kit can be used to detect ASA or T-ASA content in samples alone, or calculate DHA content after simultaneous detection of ASA and T-ASA content.
5. The samples need tested on the same day after extraction.

Experimental Examples:

1. Take 0.1g of Ziziphus jujuba Mill for sample processing, follow the measurement steps to calculate $\Delta A_{1t} = (A_t - A_c) - (A_{b1} - A_{b2}) = (0.44 - 0.019) - (0.025 - 0.009) = 0.402$, $\Delta A_{1s} = A_s - A_{b1} = 0.341 - 0.024 = 0.317$, $\Delta A_{2t} = (A_t - A_c) - (A_{b1} - A_{b2}) = (0.591 - 0.020) - (0.024 - 0.008) = 0.555$, $\Delta A_{2s} = A_s - A_{b1} = 0.375 - 0.024 = 0.351$, calculate the AsA content and T-AsA content according to the sample weight, and get:

$$\text{AsA (nmol/g weight)} = 500 \times \Delta A_{1t} \div \Delta A_{1s} \div W = 6340.7 \text{ nmol/g weight}$$

$$\text{T-AsA (nmol/g weight)} = 500 \times \Delta A_{2t} \div \Delta A_{2s} \div W = 7905.9 \text{ nmol/g weight}$$

Related Products:

NA0776/NA0535 Ascorbic Acid(AsA) Content Assay Kit

NA0775/NA0534 Dehydroascorbic Acid(DHA) Content Assay Kit

NA0773/NA0532 Ascorbic Acid Oxidase(AAO) Activity Assay Kit

NA0852/NA0610 Ascorbate Peroxidase(APX) Activity Assay Kit

NA0817/NA0575 Monodehydroascorbate Reductase(MDHAR) Activity Assay Kit

NA0816/NA0574 Dehydroascorbate Reductase(DHAR) Activity Assay Kit