

Soil α -Glucosidase (S- α -GC) Activity Assay Kit

Note: It is necessary to predict 2-3 large difference samples before the formal determination.

Operation Equipment: Spectrophotometer

Cat No: NA0645

Size: 50T/24S

Components:

Reagent I: Toluene 2 mL \times 1, store at 4°C; **Self-provided reagent;**

Reagent II: Powder \times 2, store at -20°C. Add 10 mL distilled water per bottle when the solution will be used. Mix thoroughly. The rest of reagent store at -20°C;

Reagent III: Liquid 30 mL \times 1, store at 4°C;

Reagent IV: Liquid 60 mL \times 1, store at 4°C;

Standard solution: Liquid 1 mL \times 1, 5 mmol/L of p-nitrophenol solution. Dilute the standard 50 times with reagent III before use to obtain 100 μ mol/L standard solution.

Product Description:

S- α -GC can catalyze the hydrolysis of glycosidic bonds between aryl or hydrocarbon groups and glycosyl groups to form glucose. It is one of the important components of cellulolytic enzyme system. It has important physiological functions in carbohydrate metabolism of soil microorganisms.

S-a-GC can catalyze the formation of p-nitrophenol from p-nitrobenzene-a-d-glucopyranoside. It has characteristic light absorption at 400 nm.

Required but Not Provided:

Spectrophotometer, desk centrifuge, water-bath, transferpettor, 1 mL glass cuvette, 50 mesh sieve(or smaller), toluene (express delivery is not allowed) and distilled water.

Protocol

I. Preparation:

The fresh soil sample is dried by natural air or air in a 37°C oven, and passes through 30-50 mesh sieve.

II. Determination procedure:

1. Preheat spectrophotometer for 30 min, adjust wavelength to 400 nm, set the counter to zero with distilled water.

2. Operation table:

Reagent (μ L)	Test tube (A_T)	Control tube (A_C)	Standard tube (A_S)	Blank tube (A_B)
Sample	0.1	0.1	-	-
Reagent I	25	25	-	-
Mix thoroughly. Moisten the soil sample and place it at room temperature for 15 min.				

Reagent II	400	-	-	-
Reagent III	500	500	-	-
Mix thoroughly. Water bath at 37°C for 1 h. Boiling water bath for 5 min (cover tightly to prevent water loss). Cool it with running water/ice bath.			-	-
Reagent II	-	400	-	-
Centrifugate at 10000 rpm, 25°C for 10 min. Take the supernatant for test.			-	-
Supernatant	500	500	-	-
Standard	-	-	500	-
Distilled water	-	-	-	500
Reagent IV	1000	1000	1000	1000

Mix thoroughly. Stand for 2 min at room temperature. Determine the absorption value A. Record as A_T , A_C , A_S , A_B . $\Delta A = A_T - A_C$. $\Delta A_S = A_S - A_B$. Each test tube is provided with a contrast tube. Blank tube just need test once or twice.

III. S- α -GC Calculation:

Unit definition: One unit of enzyme is defined as the amount of enzyme that catalyzes the production of 1 μmol p-nitrophenol per day every gram tissue weight in the reaction system.

$$\text{S-}\alpha\text{-GC (U/g weight)} = \Delta A \div (\Delta A_S \div C_S) \times V_{RT} \div W \div T = 2.22 \times \Delta A \div \Delta A_S \div W$$

V_{RT} : Total reaction volume, 9.25×10^{-4} L;

C_S : Concentration of standard solution, 100 $\mu\text{mol/L}$;

T: Reaction time, 1 h = 1/24 d;

W: Sample weight, g.

Note:

1. If $\Delta A < 0.01$, the water bath time of 37°C can be prolonged. If $\Delta A > 1.5$, the supernatant can be diluted before test. Pay attention to the change of various factors in the final calculation.

Experimental Example:

1. Two tubes of 0.02g clover soil are taken as the test tube and control tube. According to the test steps, measured: $\Delta A = A_T - A_C = 0.677 - 0.369 = 0.308$, $\Delta A_S = A_S - A_B = 0.702 - 0.007 = 0.695$.

S- α -GC activity (U/g soil sample) = $2.22 \times \Delta A \div \Delta A_S \div W = 2.22 \times 0.308 \div 0.695 \div 0.1 = 9.8383$ U/g soil sample.

2. Take two tubes of 0.02g forest soil sample, namely the test tube and the control tube, according to the determination steps, use 96 well plate to measure and calculate $\Delta A = A_T - A_C = 0.629 - 0.361 = 0.268$, $\Delta A_S = A_S - A_B = 0.702 - 0.007 = 0.695$.

S- α -GC activity (U/g soil sample) = $2.22 \times \Delta A \div \Delta A_S \div W = 2.22 \times 0.268 \div 0.695 \div 0.1 = 8.5606$ U/g soil sample.

Related Products:

NA0858/NA0616 Soil β -glucosidase (β -GC) Activity Assay Kit

NA0373/NA0372 Soil Neutral Invertase(S-NI) Activity Assay Kit

NA0850/NA0608 Soil Saccharase(S-SC) Activity Assay Kit