

Plant Lipxygenase (LOX) Activity Assay Kit

Note: Take two or three different samples for prediction before test.

Operation Equipment: Spectrophotometer

Catalog Number: NA0842

Size: 50T/48S

Product composition:

Extract solution: Liquid 50 mL×1. Storage at 4°C. This reagent contains insoluble substances, just mix well before use.

Reagent I: Liquid 45 mL×1. Storage at 4°C.

Reagent II: Powder×1. Storage at 4°C. Add 5 mL of Reagent I to dissolve the powder, and add 0.1 mL of 0.2 mol/L NaOH for the solution to clear.

Product Description

Lipxygenase (LOX) is widely found in plant tissues, especially soybean seeds. LOX catalyzes the oxidation of unsaturated fatty acids, resulting in membrane lipid peroxidation. It plays an important role in plant growth and development, maturation and aging.

LOX catalyzes the oxidation of linoleic acid, the oxidation product has a characteristic absorption peak at 234 nm. The rate of increase in absorbance at 234 nm is measured to calculate the LOX activity.

Reagents and Equipment Required but Not Provided.

Spectrophotometer, 1 mL quartz cuvette, refrigerated centrifuge, adjustable pipette, mortar/ homogenizer, ice and distilled water

Procedure:

I. Sample Extraction:

Tissue sample: Weigh about 0.1 g of sample and add 1 mL of Extract solution, fully grind on ice, centrifuge at 16000 ×g and 4 °C for 20 minutes, and take the supernatant for testing.

II. Determination procedure:

1. Preheat the spectrophotometer for more than 30 minutes, adjust the wavelength to 234 nm, and set zero with distilled water.
2. Reagent I is incubated in a water bath at 25°C for more than 30 minutes.
3. Blank tube: In a 1 mL quartz cuvette, add 100 μL of distilled water, 800 μL of Reagent I and 100 μL of Reagent II, after mix them quickly, measure at 234 nm, record the absorbance at 15s and 75s, and record them as A1 and A2.
4. Test tube: Add 100 μL of supernatant, 800μL of Reagent I and 100μL of Reagent II to a 1 mL quartz cuvette, after mix them quickly, measure at 234 nm, record the absorbance at 15s and 75s, and record them as A3 and A4.

III. Calculations

1. Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the absorbance of 0.001 change at 37 °C in 1 milliliter reaction system per minute every milligram protein.

$$\text{LOX activity (U/mg)} = [(A4-A3)-(A2-A1)] \div 0.001 \div (Cpr \times Vs) \div T \times Vr = 10^4 \times [(A4-A3)-(A2-A1)] \div Cpr$$

2. Sample weight:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the absorbance of 0.001 change at 37 °C in 1 milliliter reaction system per minute every gram tissue sample.

$$\text{LOX activity (U/g)} = [(A4-A3)-(A2-A1)] \div 0.001 \div (W \times Vs \div Ve) \div T \times Vr = 10^4 \times [(A4-A3)-(A2-A1)] \div W$$

Cpr: Supernatant protein concentration, mg/mL;

T: Reaction time, 1 minute;

Vs: Sample volume, 0.1 mL;

Ve: Extraction volume, 1 mL;

Vr: Reaction volume, 1 mL;

W: Sample weight, g.

Notes:

1. Reagent II is susceptible to spontaneous oxidation, resulting in a high blank tube measurement value, which must be prepared before use.
2. Sample preparing process and other processes need to be performed on ice, and the enzyme activity measurement must be completed on the same day.
3. Before the formal experiment, do 1-2 pre-experiments to ensure that ΔA is in the range of 0.02 - 1.2. If the reaction is a obvious suspension, please measure it after dilution.

References:

[1] Dou S, Liu S, Xu X, et al. Octanal inhibits spore germination of *Penicillium digitatum* involving membrane peroxidation[J]. *Protoplasma*, 2017, 254(4): 1539-1545.

Related products:

NA0822/NA0580 Free fatty acid (FFA) content detection kit

NA0701/NA0460 Lipase (LPS) activity detection kit

NA0790/NA0549 Ethanol dehydrogenase (ADH) activity detection kit

NA0791/NA0550 Pyruvate decarboxylase (PDC) activity detection kit

NA0819/NA0577 Triglyceride (TG) content detection kit

NA0733/NA0491 Free cholesterol (FC) content detection kit

NA0808/NA0566 Acetaldehyde dehydrogenase (ALDH) activity detection kit

NA0834/NA0592 Acetyl-coa carboxylase (ACC) activity detection kit

NA0727/NA0485 Total cholesterol (TC) assay kit