

# Sorbitol Dehydrogenase(SDH) Activity Assay Kit

**Note:** Take two or three different samples for prediction before test.

**Operation Equipment:** Spectrophotometer

**Cat No:** NA0689

**Size:**50T/48S

## Components:

Extract solution: 60 mL×1. Storage at 4°C.

Reagent I: Powder×1. Storage at 4°C. Add 50 mL of Reagent V into Reagent I powder, dissolve and store at 4°C.

Reagent II: Powder×1. Storage at -20°C. Before use, add 12.5 mL of Reagent V, after fully dissolved, 2.5 mL of per bottle can be sub packed and stored at -20°C;

Reagent III: Powder×1. Storage at -20°C. Before use, add 12.5 mL of Reagent V, after fully dissolved, 2.5 mL of per bottle can be sub packed and stored at -20°C;

Reagent IV: Powder×5. Storage at -20°C. Add 10 mL of Reagent I before use, which is the working solution. Prepared when the solution will be used, and it is deteriorated in 24 hours.

Reagent V: 80 mL×1. Storage at 4°C.

Standard: Powder×1. Storage at -20°C. Add 1.4 mL of distilled water before use, which is 10 μmol/mL NADH standard.

## Product Description:

SDH (EC 1.1.1.14) catalyzes the dehydrogenation of sorbitol to fructose, which is one of the key enzymes to regulate the content of sorbitol in vivo.

SDH catalyzes the dehydrogenation of sorbitol to fructose, and the reduction of NAD<sup>+</sup> to NADH. The generated NADH can transfer electrons to NBT to generate purple hairpin. According to this principle, SDH activity can be calculated.

## Reagents and Equipment Required but Not Provided:

Spectrophotometer, desk centrifuge, water bath, transferpettor, 1 mL glass cuvette, mortar/homogenizer, ice, distilled water.

## Procedure

### I. Sample preparation:

a. Bacteria or cells

Collecting bacteria/cells into the centrifuge tube. The supernatant is discarded after centrifugation. The ratio of bacteria/cell amount ( $10^4$ ): the volume of Extract solution (mL) is 500~1000:1 (it is suggested to take about 5 million bacteria/cell and add 1 mL of Extract solution). Bacteria/cell is splitted by ultrasonication(placed on ice, 200W, work time 3s, interval 10s, repeat for 30 times). Centrifuge at 8000×g for 10 minutes at 4°C, and the supernatant is used for test.

## b. Tissue

The ratio of tissue mass (g): the volume of Extract solution (mL) = 1: 5~10 (it is suggested to take about 0.1 g of tissue and add 1 mL of Extract solution), ice-bath homogenate. Centrifuge at  $8000 \times g$  for 10 minutes at  $4^{\circ}\text{C}$ , and the supernatant is used for test.

## c. Serum (plasma) sample:

Detect sample directly.

## II. Determination procedure

a. Preheat the spectrophotometer for 30 minutes, adjust wavelength to 570 nm, set zero with distilled water.

## b. standard solution

The  $10 \mu\text{mol/mL}$  NADP is respectively diluted 1, 0.9, 0.8, 0.7, 0.6, 0.5, 0.4 and  $0.3 \mu\text{mol/mL}$  NADP standard solution by distilled water. Then operate according to the following table.

Reagent name ( $\mu\text{L}$ )	Standard tube (S)	Blank tube (B)
Standard solution	100	-
Distilled water	-	100
Reagent II	150	150
Reagent III	150	150
Reagent IV	600	600

After mixing, place it at room temperature for 20 minutes, measure the absorbance of standard tube and blank tube at 570 nm respectively, record it as  $A_S$ ,  $A_B$ , calculate  $\Delta A_S = A_S - A_B$ .

## c. Sample Test

Reagent name ( $\mu\text{L}$ )	Test tube (T)
Sample	100
Reagent II	150
Reagent III	150
Reagent IV	600

Add the above reagents to the 1 mL glass cuvette in sequence, start timing at the same time of adding samples, record the initial absorbance  $A_1$  at 10 s. Put the cuvette together with the reaction solution into a water bath at  $37^{\circ}\text{C}$  (mammal) or  $25^{\circ}\text{C}$  (other species) for 3 minutes after color comparison, take out the cuvette quickly and dry it. Determinate at 570 nm, record the absorbance at 3min10s  $A_2$ , calculate  $\Delta A_T = A_1 - A_2$ .

## III. SDH Calculations

### 1. Drawing of standard curve

Take  $\Delta A_S$  as y-axis, take standard solution concentration as x-axis, draw standard curve, get standard equation  $y=kx+b$ , bring  $\Delta A_T$  into equation to get x ( $\mu\text{mol/mL}$ )

### 2. Calculate the activity of SDH

#### (1) Serum (plasma) sample SDH activity

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes the generation

of 1 nmol of NADH in the reaction system per minute every mL serum (plasma) .

$$\text{SDH (U/mL)} = 1000 \times X \times V_S \div V_{STV} \div T = 333 \times X$$

(2) Tissue, bacteria or cultured cells SDH activity

a. Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes the generation of 1 nmol of NADH in the reaction system per minute every mg protein.

$$\text{SDH (U/mg prot)} = 1000 \times X \times V_S \div (V_S \times \text{Cpr}) \div T = 333 \times X \div \text{Cpr}$$

b. Sample weight

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes the generation of 1 nmol of NADH in the reaction system per minute every g sample.

$$\text{SDH (U/g fresh wight)} = 1000 \times X \times V_S \div (W \times V_S \div V_{STV}) \div T = 333 \times X \div W$$

c. Cell amount

Unit definition: One unit of enzyme activity is defined as the amount of enzymes catalyzes the generation of 1 nmol of NADH in the reaction system per minute every  $10^4$  cells.

$$\text{SDH (U/10}^4 \text{ cell)} = 1000 \times X \times V_S \div (500 \times V_S \div V_{STV}) \div T = 0.666 \times X$$

$V_S$ : Add the volume of sample, 0.1 mL;

$V_{STV}$ : The volume of extract, 1 mL;

T: Reaction time, 3 minutes;

Cpr: The concentration of sample protein, mg/mL;

W: Sample weight, g;

500: Total number of bacteria or cells, 5 million;

$10^3$ :  $1 \mu\text{mol} = 10^3 \text{ nmol}$ .

#### **Note:**

1. When the absorbance value of spectrophotometer is greater than 0.7, it is recommended to measure after dilution.
2. Place the sample and working solution on the ice during the determination to avoid denaturation and deactivation.
3. The temperature of the reaction solution in the cuvette must be kept at 37°C or 25°C. Take a small beaker and put it into a certain amount of 37°C or 25°C distilled water. Put the beaker into a 37°C or 25°C water bath. Put the cuvette and reaction solution into the beaker during the reaction.

#### **References:**

Aguayo M F, Ampuero D, Mandujano P, et al. Sorbitol dehydrogenase is a cytosolic protein required for sorbitol metabolism in Arabidopsis thaliana[J]. Plant science, 2013, 205: 63-75.

#### **Related Products:**

- NA0695/NA0454 Plant Tissue Fructose Content Assay Kit
- NA0688/NA0447 Cellulase(CL) Activity Assay Kit
- NA0691/NA0450 Trehalase Activity Assay Kit