

Phenylalanine Ammonia-lyase (PAL) Activity Assay Kit

Note: It is necessary to predict 2-3 large difference samples before the formal determination.

Operation Equipment: Spectrophotometer

Catalog Number: NA0853

Size: 50T/48S

Components:

Extract solution: Liquid 60 mL×1. Storage at 4°C.

Reagent I: Liquid 40 mL×1. Storage at 4°C.

Reagent II: Powder×3. Storage at 4°C. Dissolve with 4 mL of distilled water before use. The reagent should be prepared just before use, and could be stored at 4°C for 2 weeks.

Reagent III: Liquid 5 mL×1. Storage at 4°C.

Product Description:

Phenylalanine Ammonia-lyase (PAL) is widely found in various plants and a few microorganisms. It is a key enzyme in phenylpropanoid metabolism. PAL is closely related to some important secondary substances synthetic such as lignin, isoflavones phytoalexin, flavonoid pigments, and play an important role in normal growth and development in plants and against the bacteria resist.

L-phenylalanine can be decomposed into trans-cinnamic acid and ammonia by PAL, and trans-cinnamic acid has the maximum absorption value at 290 nm. In this kit, the activity of PAL can be calculated by measuring the absorbance increased rate.

Reagents and Equipments Required but Not Provided:

Spectrophotometer, water bath, refrigerated centrifuge, transferpettor, 1 mL quartz cuvette, mortar /homogenizer, ice and distilled water.

Procedure:

I. Sample preparation:

Add 1 mL of Extract solution into 0.1 g of tissue, and fully homogenized on ice. Centrifuge at 10000 ×g for 10 minutes at 4°C to remove insoluble materials and take the supernatant on ice for testing.

II. Determination procedure:

1. Preheat the spectrophotometer for more than 30 minutes, adjust the wavelength to 290 nm, and set the counter to zero with distilled water.

2. Add the reagents as following

Reagent (μL)	Test tube (A1)	Contrast tube (A2)
Sample	20	-
Reagent I	780	800

Reagent II	200	200
Mix thoroughly, incubate at 30°C for 30 minutes.		
Reagent III	40	40

Mix thoroughly and place for 10 minutes. Detect the absorbance of the test tube (A1) and the contrast tube (A2) at 290 nm, calculate $A=A1-A2$.

Note: Contrast tube just needs to test once or twice.

III. Calculation:

1. Protein concentration:

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the absorbance of 0.1 change at 290 nm in per milliliter reaction system per minute every milligram protein.

$$\text{PAL(U/mg prot)} = A \times V_{rv} \div V_s \div T \div 0.1 \div \text{Cpr} = 17.3 \times A \div \text{Cpr}$$

2. Sample weight

Unit definition: One unit of enzyme activity is defined as the amount of enzyme catalyzes the absorbance of 0.1 change at 290 nm in per milliliter reaction system per minute every gram tissue.

$$\text{PAL(U/g weight)} = A \times V_{rv} \div (W \times V_s \div V) \div T \div 0.1 = 17.3 \times A \div W$$

Cpr: Sample concentration, mg/mL;

W: Sample weight, g;

Vs: Sample volume, 20 μL =0.02 mL;

V: Total sample volume, 1 mL

Vrv: Total reaction volume, 1040 μL =1.04 mL;

T: Reaction time, 30 minutes.

Recent Product Citations:

[1] Li B, Ding Y, Tang X, et al. Effect of L-Arginine on Maintaining Storage Quality of the White Button Mushroom (*Agaricus bisporus*) [J]. Food and Bioprocess Technology, 2019, 12(4): 563-574.

[2] Zhang J, Lv J, Xie J, et al. Nitrogen Source Affects the Composition of Metabolites in Pepper (*Capsicum annuum* L.) and Regulates the Synthesis of Capsaicinoids through the GOGAT–GS Pathway[J]. Foods, 2020, 9(2): 150.

References:

[1] Aydaş S B, Ozturk S, Aslım B. Phenylalanine ammonia lyase (PAL) enzyme activity and antioxidant properties of some cyanobacteria isolates[J]. Food chemistry, 2013, 136(1): 164-169.

[2] Rosler J, Krekel F, Amrhein N, et al. Maize phenylalanine ammonia-lyase has tyrosine ammonia-lyase activity[J]. Plant physiology, 1997, 113(1): 175-179.

[3] Cheng G W, Breen P J. Activity of phenylalanine ammonia-lyase (PAL) and concentrations of anthocyanins and phenolics in developing strawberry fruit[J]. Journal of the American Society for Horticultural Science, 1991, 116(5): 865-869.

Related Products:

NA0855/NA0613 Polyphenol Oxidase(PPO) Activity Assay Kit

NA0857/NA0615 Superoxide Dismutase(SOD) Activity Assay Kit

NA0854/NA0386 Catalase(CAT) Activity Assay Kit

NA0864/NA0621 Peroxidase(POD) Activity Assay Kit